

University of Groningen

Dynamics of molecular beams in a traveling-wave Stark decelerator

Zapara, Artem

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Zapara, A. (2019). *Dynamics of molecular beams in a traveling-wave Stark decelerator*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Propositions

accompanying the dissertation

Dynamics of molecular beams in a traveling-wave Stark decelerator

by

Artem Zapara

1. In order to double the effective energy scale probed by a low-energy experiment of the electron EDM, one has to quadruple the total statistics. Presently, achieving the same goal in a high-energy experiment necessitates development of new particle acceleration technologies.
2. Examination of heavy molecules with high-precision spectroscopy connects phenomena at fundamentally different scales: from the size of elementary particles (10^{-15} m) to that of the observable universe (10^{27} m).
3. For a molecule in a Stark decelerator, the internal state dynamics is largely governed by the rotation rate of the electric field.
4. The same Monte-Carlo algorithm used to approximately determine π can also be applied to calculate the phase-space acceptance of a decelerator.
5. Searching for a molecular signal in a noisy background with laser-induced fluorescence is analogous to looking for a black cat in a dark room: it is quite difficult to find, especially when there is no cat.
6. The concept of a nonadiabatic transition encompasses the mutability of processes in physics, biology and even in the social disciplines such as economics.
7. Having a good sense of smell can be vital in saving expensive laboratory equipment from total failure.
8. To satisfy a substantial public demand, every academic researcher should popularize science.